

Amylolytic enzymes from the yeast *Lipomyces kononenkoae*

Nivetha RAMACHANDRAN¹, Isak S. PRETORIUS^{1,2} & Ricardo R. CORDERO OTERO^{1*}

¹*Institute for Wine Biotechnology, Department of Viticulture & Oenology, Stellenbosch University, Victoria Street, Stellenbosch, ZA 7600, South Africa; phone: ++ 27 21 8083770, fax: ++ 27 21 8083771, e-mail: rrc@sun.ac.za*

²*The Australian Wine Research Institute, Waite Road, Urrbrae, Adelaide, SA 5064, Australia*

Abstract: The *Lipomyces kononenkoae* α -amylases LKA1 and LKA2 belong to the glycoside hydrolase family 13 and exhibit specificity towards α -1,4 and α -1,6 linkages in starch and related substrates. LKA1 exhibits specificity towards α -1,4 and α -1,6 linkages and large amounts of reducing sugars are liberated from highly branched amylopectin and glycogen and linear amylose. LKA2, on the other hand, shows high reactivity towards lintner starch, dextrin and amylose, although only small amounts of reducing sugars are liberated from branched substrates, such as amylopectin and glycogen. These enzymes share the four conserved segments of the catalytic domain found in other members of the family, but have some major variant amino acids within these segments. In addition, LKA1 consists of an N-terminal starch-binding domain (SBD). This is the only α -amylase known to possess this N-terminal domain and it exhibits homology to the N-terminal SBD of *Rhizopus oryzae* glucoamylase. It shares no homology with the C-terminal starch-binding domains present in the cyclodextrin glucanotransferases, glucoamylases or α -amylases. The evolutionary tree based on the sequence alignment of SBDs reveals that the N-terminal SBDs are separated from the C-terminal SBDs.

Key words: *Lipomyces kononenkoae* α -amylases, sequence analyses, substrate specificity.

Abbreviations: CGTase, cyclodextrin glucanotransferase; GH, glycoside hydrolase; LKA1, *Lipomyces kononenkoae* α -amylase 1; LKA2, *Lipomyces kononenkoae* α -amylase 2; SBD, starch-binding domain.

Introduction

Starch is a major renewable resource consisting of two types of molecules, amylose, which generally makes up 20–30%, and amylopectin, which generally constitutes 70–80%. Both these types of molecules consist of polymers of α -D-glucose units in the ⁴C₁ conformation. In amylose, these are linked α -(1,4), with the ring oxygen atoms all on the same side, whereas in amylopectin, about one residue in every twenty or so is also linked α -(1,6), forming branch points (IMMEL et al., 2000). The use of amylolytic enzymes to break down starch to soluble sugars has become increasingly important in the production of various commercial products, such as fuel ethanol, alcoholic beverages and sweeteners. α -Amylases were originally recognised as a group of starch hydrolases and related enzymes that exhibit clear sequence similarities and a predicted common super secondary fold, a parallel (β/α)₈ barrel (FARBER & PETSKO, 1990; FARBER, 1993). The α -amylase family is the largest glycoside hydrolase (GH) family comprising over 30 different specificities from hydrolases, transferases and isomerases. Currently, the α -amylase family con-

stitutes the clan GH-H of families 13, 70 and 77, and the enzymes of these different families can operate on α -1,1-, α -1,2-, α -1,3- and α -1,5-linkages, as well as on α -1,4- and α -1,6-glucosidic linkages (VAN DER MAAREL et al., 2002). In nature, there are more than 150 starch-assimilating yeasts, but only a few of them secrete a combination of enzymes that can cleave the α -1,4 and α -1,6 linkages of the complex starch molecule. Several yeasts, such as *Saccharomycopsis fibuligera* (ITO et al., 1987), *Shizosaccharomyces pombe* (WOOD et al., 2002), *Cryptococcus neoformans* (IEFUJI et al., 1996) produce α -amylases; the yeast species *Schwanniomyces occidentalis* secretes two α -amylases SWA1 and SWA2, which were expressed in *Saccharomyces cerevisiae*. SWA1 and SWA2 secreted from *S. cerevisiae* showed a significant modification of their properties when these were compared with those of the wild-type enzymes (ABARCA et al., 1989; 1991; YANEZ et al., 1998). The ability of the yeast species *Lipomyces kononenkoae* to degrade raw starch completely has been well documented (SPENCER MARTINS & VAN UDEN, 1979; WILSON & INGLEDEW, 1982; DE MOT et al., 1984) SPENCER MARTINS & VAN UDEN (1979) eval-

* Corresponding author

uated 81 raw starch-assimilating yeasts, representing 59 species, and reported that the highest biomass production on starch media was obtained with strains of *Lipomyces kononenkoae* and *Lipomyces starkeyi*. These two yeast species secrete a set of starch-hydrolysing enzymes of which the genes encoding for some of these have been cloned and expressed in simpler systems, such as *Saccharomyces cerevisiae* and *Escherichia coli* (STEYN & PRETORIUS, 1995; BIGNELL et al., 2000; EKSTEEN et al., 2003a; KANG et al., 2004). The genes encoding two novel α -amylases, LKA1 and LKA2, were isolated from *L. kononenkoae* strain IGC4052B, cloned and expressed in *S. cerevisiae* (STEYN & PRETORIUS, 1995; EKSTEEN et al., 2003b). These α -amylases exhibit different substrate specificities towards starch and related substrates, acting on their α -1,4 and α -1,6 linkages. In this review, we summarise the sequence analyses and properties of *L. kononenkoae* α -amylases LKA1 and LKA2.

Amylolytic system of *Lipomyces kononenkoae*

All the starch-assimilating yeasts necessarily produce one or more extracellular amylolytic enzymes. The crude concentrates of culture fluids of *Lipomyces* strains are able to extensively hydrolyse a wide variety of polysaccharides, including starch, and two strains of *L. kononenkoae* were identified to produce a different extracellular amylolytic system capable of total starch hydrolysis (SPENCER MARTINS & VAN UDEN, 1979). However, due to their complex genetics, only a few genes encoding these enzymes have been identified and further characterised. The culture filtrate from *L. kononenkoae* strain CBS5608 consists of three different enzymes: α -amylase, glucoamylase, and a third enzyme with both debranching activity on amylopectin and transferase activity hydrolysing α -1,6 chains in panose. *L. kononenkoae* strain IGC4051 was reported to secrete an extracellular isoamylase, which increases the amyolysis of amylopectin and glycogen, completely hydrolysing these substrates into maltose when combined with a β -amylase. This enzyme does not have any action on pullulan or dextrin (SPENCER MARTINS, 1982).

The strain *L. kononenkoae* IGC4052B, which is a catabolite-derepressed mutant of IGC4052, secretes two enzymes with activity on starch, namely a novel raw starch-degrading α -amylase, LKA1, and a second α -amylase, LKA2. LKA1 α -amylase acts on glucose polymers containing α -1,4 and α -1,6 bonds by endohydrolysis, producing maltose, maltotriose and maltotetraose. LKA2 exhibits properties of α -amylase, but also possesses side activity towards dextrin. The genes encoding these enzymes have been identified, cloned and expressed in *Saccharomyces cerevisiae* (STEYN & PRETORIUS, 1995; EKSTEEN et al., 2003a). The expression of LKA1 in *S. cerevisiae* in conjunction with LKA2 results in higher ethanol yields as a result of the synergistic action of these enzymes (EKSTEEN et al., 2003b).

Biochemical properties of *L. kononenkoae* α -amylases

LKA1 α -amylase has a theoretical molecular mass of approximately 76 kDa (STEYN & PRETORIUS, 1995). The enzyme exhibited properties of a glycoprotein, indicated by DIF-glycan detection. The temperature optimum of LKA1 ranged from 40 to 50°C (STEYN & PRETORIUS, 1995); and this is within the range of optimum temperatures reported for other yeast α -amylases (DE MOT, 1990; IEFUJI et al., 1996). The stability of the enzyme is maintained between 20 and 50°C, however, and it retains 83% of its activity at 55°C. The optimal pH range of 3-8 was similar to the values reported for *L. starkeyi* α -amylase (KELLY et al., 1985) and other yeast amylases (DE MOT, 1990).

LKA1 α -amylase activity was partially inhibited by Cu^{2+} and Zn^{2+} , whereas other metal ions, such as Co^{2+} and Mg^{2+} , had a slight stimulating effect. Calcium, which stabilises most α -amylases, had no effect on the activity of LKA1. Enzymatic activity was not affected by EDTA or EGTA, but was stimulated by SDS and low concentrations of urea, and was sensitive to acetone.

The second α -amylase expressed in *S. cerevisiae*, LKA2, had a temperature optimum of 60°C and maintained its maximum activity in the temperature range of 50-65°C. The optimum pH for activity was estimated to be 3.5. The molecular weight of this protein was around 55 kDa and, similar to the LKA1 α -amylase, neither Ca^{2+} nor EDTA had any effect on enzymatic activity (STEYN & PRETORIUS, 1995; EKSTEEN et al., 2003a).

LKA1 exhibited specificity towards α -1,4 and α -1,6 linkages and large amounts of reducing sugars were liberated from the highly branched amylopectin and glycogen and amylose. The k_m values for lintner starch and amylopectin were 2.31 and 1.97 mg/mL respectively. LKA1 also exhibited a side activity on pullulan. LKA2, however, showed high reactivity towards lintner starch, dextrin and amylose, but only small amounts of reducing sugars were liberated from branched substrates, such as amylopectin and glycogen (EKSTEEN et al., 2003a). The relative activities of the LKA1 and

Table 1. Activity of *Lipomyces kononenkoae* α -amylases on starch and related substrates (STEYN & PRETORIUS, 1995; EKSTEEN et al., 2003a).

| Glucose polymer | Relative enzyme activity (%) | |
|----------------------|------------------------------|------|
| | LKA1 | LKA2 |
| Amylopectin (potato) | 100 | 17 |
| Starch (lintner) | 95 | 100 |
| Glycogen (oyster) | 75 | 9 |
| Dextrin | 65 | 91 |
| Pullulan | 2 | 2 |

Table 2. Variations between and similarity within the conserved regions of the catalytic domain of *Lipomyces kononenkoae* amylases.

| Enzymes | Source | Region 1 | Region 2 | Region 3 | Region 4 |
|-------------------|--|------------|----------------|----------|------------|
| α -Amylase | <i>Aspergillus oryzae</i> | 117 DVVANH | 202 GLRIDTVKH | 230 EVLD | 292 FVENHD |
| Pullulanase | <i>Klebsiella aerogenes</i> | 600 DVVYNH | 671 GFRFDLMGY | 704 EGWD | 827 YVSKHD |
| Neopullulanase | <i>Bacillus stearothermophilus</i> | 242 DAVFNH | 324 GWRDLVANE | 357 EIWH | 419 LLGSHD |
| Amylopullulanase | <i>Clostridium thermohydrosulfuricum</i> | 488 DGVFNH | 594 GWRDLVANE | 627 ENWN | 699 LLGSHD |
| CGTase | <i>Bacillus macerans</i> | 135 DFAPNH | 225 GIRFDVAVKH | 258 EWFL | 324 FIDNHD |
| Amylomaltase | <i>Streptococcus pneumoniae</i> | 224 DMWAND | 291 IVRIDHFRG | 332 EELG | 391 YTGTHD |
| Isoamylase | <i>Pseudomonas amyloclavata</i> | 292 DVVYNH | 371 GFRFDLASV | 435 EPWA | 505 FIDVHD |
| LKA1 | <i>Lipomyces kononenkoae</i> | 264 DIVVNH | 248 GLRIDTVKH | 376 EVFD | 439 FLENQD |
| LKA2 | <i>Lipomyces kononenkoae</i> | 137 DVVINH | 223 GIRLDVAVKH | 251 EALN | 312 FLDNQD |
| LSA | <i>Lipomyces starkeyi</i> | 287 DIVVNH | 371 GLRIDTVKH | 400 EVFD | 462 FLENQD |

LKA2 enzymes on different glucose polymers are summarised in Table 1.

Sequence alignment of *L. kononenkoae* α -amylases

The analysis of sequences of LKA1 and LKA2 reveals homology to various yeast and plant α -amylases, namely to the α -amylases of *Aspergillus nidulans*, *Debaryomyces occidentalis*, *Saccharomycopsis fibuligera* and *Schizosaccharomyces pombe*, cyclodextrin glucanotransferases (CGTases), pullulanases, α -glucosidase, and β -amylase from *Bacillus polymyxa* (TAKIZAWA & MUROOKA, 1985; KAWAZU et al., 1987; KANEKO et al., 1998). The sequence comparison of α -amylases from different yeasts is presented in Figure 1 (EKSTEEN et al., 2003a). LKA1 α -amylase exhibits a high homology of around 78% to the α -amylase from *L. starkeyi*, whereas LKA2 has a homology of about 36% to it (KANG et al., 2004). The three-dimensional structure of *Aspergillus oryzae* TAKA amylase (MATSUURA et al., 1984) was used as a reference model for the structure interpretation of both the α -amylases, LKA1 and LKA2. The analyses of primary structures of LKA1 and LKA2 with that of TAKA amylase revealed corresponding similarities for the catalytic domain and the C-terminal stabilising domain. The sequence comparison suggested that the amino acid sequences of both the LKA1 and LKA2 included the four conserved regions comprising the strands β_3 , β_4 , β_5 , and β_7 , containing the catalytically important residues His122, Asp206, Glu230, and Asp297 respectively (HENRISSAT, 1991; TAKATA et al., 1992; SVENSSON, 1994), as previously identified in the α -amylase family of enzymes (KURIKI et al., 1991). Further, six amino acid residues, which are strictly conserved in the four conserved regions of the α -amylase family, are also completely preserved in LKA1 and LKA2. It has been suggested earlier (JANECEK, 1997) that conservations in the catalytic regions of α -amylases might be related to the maintenance of structure and variations within the segments may be characteristic of certain enzyme specificities.

Variations in the conserved segments of the catalytic domain

The sequence analyses showed some amino acid residues that are variable in the conserved segments of *L. kononenkoae* α -amylases, as shown in Table 2. Region IV of LKA1, LKA2 and LSA contains an uncharged Gln in the place of the highly conserved, charged His, which is involved in substrate binding in other α -amylases. The importance of the histidine has been well demonstrated through structural, kinetic and mutagenesis approaches. The mutation of His296 to Asn led to a 100-fold decrease in the K_{cat} values in barley α -amylase (SØGAARD et al., 1993); a similar mutation led to a loss of 50% of the original activity in CGTase from a *Bacillus*, and similar results have been observed for human pancreatic α -amylase (ISHIKAWA et al., 1993). In *Bacillus stearothermophilus* neopullulanase, the mutation of the corresponding His to Glu led to a 70% decrease in its activity towards α -1,4 linkages, leading to a lower maltose release and higher panose production (KURIKI et al., 1991). Inhibitor-binding studies conducted on barley α -amylase suggest that the imidazole group of the His residue forms a hydrogen bond with the OH of the glycone residue at the bond cleavage site. Thus, the mutation of this His296 greatly diminishes the ability of the enzyme to stabilise the transition state, as indicated by the decreases in K_{cat}/K_m for the substrates used (SØGAARD et al., 1993). These research findings reiterate the importance of the invariant histidine (corresponding with His296 of the barley α -amylase) in this family of enzymes.

The uncharged Gln in the place of His296 in the fourth conserved sequence region is unique to the *Lipomyces* α -amylases cloned and characterised thus far. This is clear from the sequence comparison of these enzymes with other members of the α -amylase family. The significance of Gln at this position in LKA1 α -amylase is currently under investigation (N. RAMACHANDRAN et al., unpublished data). It is worth mentioning that understanding the role of variant amino acids in such highly conserved regions of these enzymes might broaden our knowledge on the charac-

```

LKA1  ESVTGSNHVQLASYEMCGSTLSASLYVYNDYDKIVTLTYLTSSG---TTG-ST 50
LSA   ESVTSSNHVQLASHEMCDSTLSASLYIYNDYDKIVTLTYLTSSG---TTG-SV 50
Rg    ASIPISSASVQLDSYNYDGFSTFGKLYVKNIAYSKKVTVIYADGSDNWNNGNTI 54
Aa    NSPPDDKAVALSSYSYCGGYSASAFVKNLSYDKLVTLTYWTNADNKSTPLNAGS 54
      *      * * * :      : *      * * * * :
LKA1  LALILPVWSN-NWELWTL--SAIAAGAVEITGASYVSDTSVTYTTSS----- 94
LSA   TASYSSLSLN-NWELWTL--SAPAADAVEITGASYVSDASATYATSFDIPL- 99
Rg    AASYSAPISGSNYEYWTF--SASINGIKEF---YIKYEVSG----- 90
Aa    LDYVKAASDDQSWELWSLNVTTVPDGDALLNITYVAASIGKTNSQQLNVQVE 107
      : * * : :      :      * :

```

Fig. 2. Sequence alignment of the N-terminal SBDs of LKA1 (*Lipomyces kononenkoae* α -amylase; STEYN et al., 1995) and Rg (*Rhizopus oryzae* glucoamylase; ASHIKARI et al., 1986) with the N-terminal regions of Aa (*Arxula adeninivorans* glucoamylase; BUI et al., 1996) and LSA (*Lipomyces starkeyi* α -amylase; KANG et al., 2004). Asterisks and colons signify the identical and similar residues, respectively.

teristic properties of the individual enzymes. The significance of variations in the *LKA2*-encoded α -amylase: Leu in region 2, Ala and Asn in region 3, and Asp and Gln in region 4, is yet to be explored. In addition, *LKA2* α -amylase contains an Arg residue in the place of the Lys121 (TAKA amylase numbering), which is well conserved in the α -amylase family. Previous reports on mutational analyses of Lys to Arg/Asn in *Saccharomyces fibuligera* showed a three-fold increase in activity towards shorter substrates (MATSUI et al., 1992).

Starch binding and raw starch hydrolysis

In some distinct members, the catalytic domain is preceded by an extra sequence, a domain whose role is uncertain (MACGREGOR et al., 2001). The N-terminal region of the *LKA1* α -amylase preceding the catalytic domain consists of an extension of 132 amino acid residues that exhibits significant homology to the N-terminal regions of *L. starkeyi* α -amylase, and 36% similarity to those of the glucoamylases from *Rhizopus oryzae* and *Arxula adeninivorans* (Fig. 2). Predictions of a hypothetical secondary structure using the predict protein tools (<http://www.expasy.ch/>) suggest the presence of six β -strands. This region in *R. oryzae* has been reported to be the starch-binding domain (SBD) (ASHIKARI et al., 1986). Deletion of the N-terminal region (1-132) in *LKA1* resulted in a loss of the binding function of this enzyme to raw starch (N. RAMACHANDRAN et al., manuscript submitted). It is also known, that the presence of SBD may confer thermostability properties to enzymes (IEFUJI et al., 1996). *LKA2* α -amylase, on the other hand, possesses a short extension of 19 amino acid residues, which does not exhibit any significant homology to any known α -amylase sequences. Whether this region functions as an extension of the catalytic domain or is involved in any protein function is yet to be understood.

On the basis of the knowledge of different SBDs present in this family, they have been classified into the carbohydrate-binding module (CBM) families 20, 21, 25, 26, 34 and 41 (COUTINHO & HENRISSAT, 1999; RODRIGUEZ-SANOJA et al., 2005). In particular, the SBDs from lactobacilli classified as CBM26 have been described by SANTIAGO et al. (2005). The SBDs present at the C-terminal end of the enzymes have been classified under CBM20, along with the SBDs from CGTases (EC 2.4.1.19, domain E), α -amylases (EC 3.2.1.1), β -amylases (EC 3.2.1.2), maltotetraose-forming exo-

amylases (EC 3.2.1.60), maltogenic α -amylases (EC 3.2.1.133) and other hydrolases. The tertiary structures of nine of the family CBM20 SBDs are known thus far (CORNETT et al., 2003). The *Rhizopus oryzae* and *Arxula adeninivorans* SBDs are currently classified as the family CBM21. The sequence homology of *LKA1* to the other N-terminal SBDs is shown in Figure 2. However, no three-dimensional structures are available for this family of SBDs, which limits the knowledge of the interactions of the SBD with the other domains on a structural level. It is evident, however, that the N-terminal SBDs from *R. oryzae* glucoamylase and *LKA1* α -amylase are very distantly related to all the other known C-terminal SBDs. The differences in the sequences of the C- and N-terminal SBDs belonging to CBM20 and CBM21, respectively, may reflect the possibility that, during their evolution, the *Rhizopus* and *Aspergillus* glucoamylases obtained their SBDs independently (TANAKA et al., 1986). While most amylases possess their binding domains at the C terminus, it can be speculated that multiple domain organisations existed in ancestral fungi and that these enzymes did not follow phylogenetic lines. This might explain why *R. oryzae* and *A. adeninivorans* possess a putatively newer type of SBD at the N terminus, even though Zygomycetes like *Rhizopus* diverged from *Ascomycetes* before the appearance of yeasts (COUTINHO & REILLY, 1997). Moreover, the presence of SBD in an amylolytic enzyme is closely connected to the enzyme origin (JANECEK et al., 2003). The presence of the N-terminal region in the *LKA1*-encoded α -amylase homologous to the SBD of *R. oryzae* glucoamylases might suggest the independent evolutionary behaviour of SBDs in glycoside hydrolases. BORK et al. (1998) further reported alignment of the SBDs of bacterial and fungal amylases and a bacterial CGTase with some mammalian glycogen-targeting subunits. Strikingly, the amino acids Gly146, Asn152, Lys157, Asp189, Phe193, Tyr213 and Asn224 of these mammalian polysaccharide-binding domains are well-conserved only in the N-terminal SBD of *R. oryzae* and *LKA1*. No similarity was noted in any of the C-terminal SBDs examined.

The differences between the C-terminal SBDs and the N-terminal SBDs are clearly reflected in their evolutionary tree (Fig. 3), which was calculated using the amino acid sequences (Table 3) according to JANECEK et al. (2003); in the current study we have included the *Lipomyces* amylases and enzymes with N-terminal starch SBDs to this evolutionary tree. The C-




Fig. 3. Evolutionary tree of SBDs of α -amylases, glucoamylases and CGTases.




Fig. 4. Location of the starch binding domains in glucosyl hydrolase members. **CD**, catalytic domain; **SBD**, starch binding domain; **C**, domain C; **D**, domain D.

terminal SBDs of α -amylases and glucoamylases cluster together, while the CGTases are on the branch adjacent to the α -amylase-glucoamylase group. The *Lipomyces* enzymes, LKA1 and LSA α -amylases, are on the same branch as that of the *R. oryzae* glucoamylase SBD. The fact that the SBD occurs in GH13, GH14

and GH15 (JANECEK & SEVCIK, 1999) supports the idea that there has been a separate evolution of this domain. The different organisations found in the α -amylase family (JESPERSEN et al., 1991; JUGE et al., 2002) with regard to the SBDs are illustrated in Figure 4.

Table 3. Amino acid sequences used for comparative analyses and construction of evolutionary tree.^a

| Enzyme | Source | Abbr. | SwissProt | Reference |
|----------------------|--|---------|-----------|-----------------------------------|
| C-terminal SBDs | | | | |
| α-Amylases | <i>Aspergillus nidulans</i> | Aspnd | Q9UV09 | Unpublished |
| | <i>Aspergillus kawachii</i> | Aspka | P13296 | KANEKO et al. (1996) |
| | <i>Cryptococcus</i> sp. S2 | Crcsp | Q92394 | IEFUJI et al. (1996) |
| | <i>Streptomyces albidoflavus</i> | Stral | P09794 | LONG et al. (1987) |
| | <i>Streptomyces griseus</i> | Strgr | P30270 | VIGAL et al. (1991) |
| | <i>Streptomyces lividans</i> TK21 | Strli21 | O86876 | Unpublished |
| | <i>Thermomonospora curvata</i> | Thscu | P29750 | PETRICEK et al. (1992) |
| Maltotetraohydrolase | <i>Pseudomonas saccharophila</i> | Psesa | P22963 | ZHOU et al. (1989) |
| Maltopentaohydrolase | <i>Pseudomonas</i> sp.KO-8940 | Psesp | Q52516 | SHIDA et al. (1992) |
| Maltogenic α-amylase | <i>Bacillus stearothermophilus</i> | Bacst | P19531 | DIDERICHSEN & CHRISTIANSEN (1988) |
| CGTases | <i>Bacillus</i> sp. 1-1 | Bac11 | P31746 | SCHMID et al. (1988) |
| | <i>Bacillus</i> sp. 17-1 | Bac17 | P30921 | KANEKO et al. (1989) |
| | <i>Bacillus</i> sp.38-2 | Bac38 | P09121 | KANEKO et al. (1988) |
| | <i>Bacillus</i> sp.6.6.3 | Bac663 | P31747 | Unpublished |
| | <i>Bacillus licheniformis</i> | Bac1i | P14014 | HILL et al. (1990) |
| | <i>Bacillus stearothermophilus</i> No 2 | Bacst2 | P31797 | FUJIWARA et al. (1992) |
| | <i>Thermoanaerobacter thermosulfurogenes</i> | Thbth | P26827 | BAHL et al. (1991) |
| | <i>Thermococcus</i> sp. B1001 | Thcsp | Q9UWN2 | YAMOMOTO et al. (1989) |
| | Glucoamylases | | | |
| | <i>Aspergillus</i> sp. X-100 | Ax | Q12537 | Unpublished |
| | <i>Aspergillus kawachii</i> | Ak | P23176 | HAYASHIDA et al. (1989) |
| | <i>Aspergillus niger</i> | An | P04064 | BOEL et al. (1984) |
| | <i>Aspergillus oryzae</i> | Ao | P36914 | HATA et al. (1991) |
| | <i>Neurospora crassa</i> | Nc | P14804 | STONE et al. (1993) |
| | <i>Humicola grisea</i> | Hg | Q12623 | Unpublished |
| | <i>Hormoconis resiniae</i> | Ho | Q03045 | JOUTSJOKEI et al. (1992) |
| | <i>Corticium rolfsii</i> | Cr | Q12596 | NAGASAKA et al. (1995) |
| N-terminal SBDs | | | | |
| α-Amylases | <i>Lipomyces kononenkoae</i> | LKA1 | Q01117 | STEYN et al. (1995) |
| | <i>Lipomyces starkeyi</i> | LSA | Q6YF33 | KANG et al. (2004) |
| Glucoamylases | <i>Arxula adeninivorans</i> | Aa | P42042 | BUI et al. (1996) |
| | <i>Rhizopus oryzae</i> | Rg | P07683 | ASHIKARI et al. (1986) |

^a Based on the study by JANECEK et al. (2003).

Future prospects

The *L. kononenkoae* α-amylases and their unique specificity properties support the rationale for structure-function relationship studies in the α-amylase family. The unique modular organisation of LKA1 also invites the construction of novel enzymes using *L. kononenkoae* α-amylases with advantageous combinations of properties, such as raw starch adsorption and activity towards a wide range of substrates. Furthermore, such studies would provide insight into various structure-function implications and would assist in unravelling how the extra domains interact with the catalytic domain or how the variant amino acids influence catalysis and specificity towards different substrates. The positioning of the N-terminal SBD, insight into cooperation between the SBD and the catalytic domain, and the investigation of the reaction mechanism and substrate specificity are among the major questions that need to be addressed in relation to the basic knowledge about these proteins.

Acknowledgements

The authors would like to thank the National Research Foundation, South Africa for funding.

References

- ABARCA, D., FERNANDEZ-LOBATO, M., CLAROS, M.G & JIMINEZ, A. 1989. FEBS Lett. **255**: 455–459.
- ABARCA, D., FERNANDEZ-LOBATO, M., DEL POZO, L & JIMINEZ, A. 1991. FEBS Lett. **279**: 41–44.
- ASHIKARI, T., NAKAMURA, N., TANAKA, Y., KIUCHI, N., SHIBONO, Y., TANAKA, T., AMACHI, T. & YOSHIZUMI, H. 1986. Agric. Biol. Chem. **50**: 957–964.
- BAHL, H., BURCHHARDT, G., SPREINAT, A., HAECKEL, K., WIENECKE, A., SCHMIDT, B. & ANTRANIKIAN, G. 1991. Appl. Environ. Microbiol. **57**: 1554–1559.
- BIGNELL, G.R., BRUCE, I.J & EVANS, I.H. 2000. Biotechnol. Lett. **22**: 1713–1718.
- BOEL, E., HANSEN, M.T., HJORT, I., HOEGH, I & FILL, N.P. 1984. EMBO J. **3**: 1581–1585.
- BORK, P., DANDEKAR, T., EISENHABER, F. & HUYNEN, M. 1998. J. Mol. Med. **76**: 77–79.
- BUI MINH, D., KUNZE, I., FOERSTER, S., WARTMANN, T., HORSTMANN, C., MANTEUFFEL, R & KUNZE, G. 1996. Appl. Microbiol. Biotechnol. **44**: 610–619.

- CLAROS, M.G., ABARCA, D., FERNANDEZ-LOBATO, M. & JIMENEZ, A. 1993. *Curr. Genet.* **24**: 75–83.
- CORNETT, C.A.G., FANG, T.Y., REILLY, P.J. & FORD, C. 2003. *Protein Eng.* **16**: 521–529.
- COUTINHO, P.M. & HENRISSAT, B. 1999. pp. 15–23. In: OHMIYA, K., HAYASHI, K., SAKKA, K., KARITA, S. & KIMURA, T. (eds) *Genetics, Biochemistry and Ecology of Cellulose Degradation*, Uni Publishers Co., Tokyo.
- COUTINHO, P.M. & REILLY, P.J. 1997. *Proteins: Struct. Funct. Genet.* **29**: 334–347.
- DE MOT, R., VAN OUDENDIJK, E. & VERACHTERT, H. 1984. *Biotechnol. Lett.* **6**: 581–586.
- DE MOT, R. & VERACHTERT, H. 1987. *Eur. J. Biochem.* **164**: 643–654.
- DE MOT, R. 1990. pp. 163–222. In: VERACHTERT, H. & DE MOT, R. (eds) *Yeast, Biotechnology and Biocatalysis*, Marcel Dekker, New York.
- DIDERICHSEN, B. & CHRISTIANSEN, L. 1988. *FEMS Microbiol. Lett.* **56**: 53–60.
- EKSTEEN, J.M., STEYN, A.J.C., VAN RENSBURG, P., CORDERO OTERO, R.R. & PRETORIUS, I.S. 2003a. *Yeast* **20**: 69–78.
- EKSTEEN, J.M., VAN RENSBURG, P., CORDERO OTERO, R.R. & PRETORIUS, I.S. 2003b. *Biotechnol. Bioeng.* **84**: 639–646.
- FARBER, G.K. 1993. *Curr. Opin. Struct. Biol.* **3**: 409–412.
- FARBER, G.K. & PETSKO, G.A. 1990. *Trends Biochem. Sci.* **15**: 228–234.
- FUJIWARA, S., KANEMOTO, M., KIM, B., LEJEUNE, A., SAKAGUCHI, K. & IMANAKA, T. 1992. *Appl. Environ. Microbiol.* **58**: 4016–4025.
- HATA, Y., KITAMOTO, K., GOMI, K., KUMAGAI, C., TAMURA, G. & HARA, S. 1991. *Agric. Biol. Chem.* **55**: 941–949.
- HAYASHIDA, S., KURODA, K., OHTA, K., KUHARA, S., FUKUDA, K. & SAKAKI, Y. 1989. *Agric. Biol. Chem.* **53**: 923–929.
- HENRISSAT, B. 1991. *Biochem. J.* **280**: 309–316.
- HILL, D.E., ALDAPE, R. & ROZZELL, J.D. 1990. *Nucleic Acids Res.* **18**: 199.
- IEFUJI, H., CHINO, M., KATO, M. & LIMURA, Y. 1996. *Biochem. J.* **318**: 989–996.
- IMMEL, S. & LICHTENTHALER, F.W. 2000. *Starch/Staerke* **52**: 1–8.
- ISHIKAWA, K., MATSUI, I., KOBAYASHI, S., NAKATANI, H. & HONDA, K. 1993. *Biochemistry* **32**: 6259–6365.
- ITO, T., YAMASHITA, I. & FUKUI, S. 1987. *FEBS Lett.* **219**: 339–342.
- JANECEK, S. 1997. *Progr. Biophys. Mol. Biol.* **67**: 67–69.
- JANECEK, S. & SEVCIK, J. 1999. *FEBS Lett.* **456**: 119–125.
- JANECEK, S., SVENSSON, B. & MACGREGOR, E.A. 2003. *Eur. J. Biochem.* **270**: 635–645.
- JESPERSEN, H.M., MACGREGOR, E.A., SIERKS, M.R. & SVENSSON, B. 1991. *Biochem. J.* **280**: 51–55.
- JOUTSJKI, V.V. & TORRKELI, T.K. 1992. *FEMS Microbiol. Lett.* **78**: 237–243.
- JUGE, N., LE GAL-COEFFET, M.F., FURNISS, C.S.M., GUNNING, A.P., KRAMHRFT, B., MORRIS, V.J., WILLIAMSON, G. & SVENSSON, B. 2002. *Biologia, Bratislava* **57 (Suppl. 11)**: 239–245.
- KANEKO, A., SUDO, S., SAKAMOTO, Y., TAMURA, G., ISHIKAWA, T. & OHBA, T. 1996. *J. Ferment. Bioeng.* **81**: 292–298.
- KANEKO, T., HAMAMOTO, T. & HORIKOSHI, K. 1988. *J. Gen. Microbiol.* **134**: 97–105.
- KANEKO, T., SONG, K.B., HAMAMOTO, T., KUDO, T. & HORIKOSHI, K. 1989. *J. Gen. Microbiol.* **135**: 3447–3457.
- KANG, H.K., LEE, J.H., KIM, D., DAY, D.F., ROBYT, J.F., PARK, K.H. & MOON, T. 2004. *FEMS Microbiol. Lett.* **233**: 53–64.
- KAWAZU, T., NAKANISHI, Y., UOZUMI, N., SASAKI, T., YAMAGATA, H., TSUKAGOSHI, N. & UDAKA, S. 1987. *J. Bacteriol.* **169**: 1564–1570.
- KELLY, C.T., MORIARTY, M.E. & FOGARTY, W.M. 1985. *Appl. Microbiol. Biotechnol.* **22**: 352–358.
- KURIKI, T., TAKATA, H., OKADA, S. & IMANAKA, T. 1991. *J. Bacteriol.* **173**: 6147–6152.
- LONG, C.M., VIROLLE, M.J., CHANG, S.Y., CHANG, S. & BIBB, M.J. 1987. *J. Bacteriol.* **169**: 5745–5754.
- MACGREGOR, E.A., JANECEK, S. & SVENSSON, B. 2001. *Biochim. Biophys. Acta* **1546**: 1–20.
- MATSUI, I., ISHIKAWA, K., MIYARI, S., FUKUI, S. & HONDA, K. 1992. *FEBS Lett.* **310**: 216–218.
- MATSUURA, Y., KUSUNOKI, W., HARADA, W. & KAKUDO, M. 1984. *J. Biochem.* **95**: 697–702.
- NAGASAKA, Y., MURAKI, N., KIMURA, A., SUTO, M., YOKOTA, A. & TOMITA, F. 1995. *Appl. Microbiol. Biotechnol.* **44**: 451–458.
- ONISHI, H. & HIDAKA, O. 1978. *Can. J. Microbiol.* **24**: 1017–1023.
- PETRICEK, M., TICHY, P. & KUNCOVA, M. 1992. *Gene* **112**: 77–83.
- RODRIGUEZ-SANOJA, R., OVIEDO, N. & SANCHEZ, S. 2005. *Curr. Opin. Microbiol.* **8**: 260–267.
- SANTIAGO, M., LINARES, L., SANCHEZ, S. & RODRIGUEZ-SANOJA, R. 2005. *Biologia, Bratislava* **60 (Suppl. 16)**: 111–114.
- SCHMID, G., ENGBRECHT, A. & SCHMID, D. 1988. pp. 71–76. In: HUBER, O. & SZEJTLI, J. (eds) *Proceedings of the Fourth International Symposium on Cyclodextrins*, Kluwer Academic Publishers, Dordrecht, Germany.
- SHIDA, O., TAKANO, T., TAKAGI, H., KADOWAKI, K. & KOBAYASHI, S. 1992. *Biosci. Biotech. Biochem.* **56**: 76–80.
- SOGAARD, M., KADZIOLA, A., HASER, R. & SVENSSON, B. 1993. *J. Biol. Chem.* **268**: 1391–1396.
- SPENCER MARTINS, I. 1982. *Appl. Environ. Microbiol.* **4**: 29–35.
- SPENCER MARTINS, I. & VAN UDEN, N. 1979. *Eur. J. Appl. Microbiol.* **6**: 241–250.
- STEYN, A.J.C. & PRETORIUS, I.S. 1995. *Curr. Genet.* **28**: 526–533.
- STONE, P.J., MAKOFF, A.J., PARISH, J.H. & RADFORD, A. 1993. *Curr. Genet.* **24**: 205–211.
- SVENSSON, B. 1994. *Plant Mol. Biol.* **25**: 141–157.
- TAKATA, H., KURIKI, T., OKADA, S., TAKESADA, Y., LIZUKA, M., MINAMIURA, N. & IMANAKA, T. 1992. *J. Biol. Chem.* **267**: 18447–18452.
- TAKIZAWA, N. & MUROOKA, Y. 1985. **49**: 294–298.
- TANAKA, Y., ASHIKARI, T., NAKAMURA, N., KIUCHI, N., SHIBANO, Y., AMACHI, T. & YOSHIZUMI, H. 1986. *Agric. Biol. Chem.* **50**: 965–969.
- VAN DER MAAREL, M.J., VAN DER VEEN, B., UITDEHAAG, J.C., LEEMHUIS, H. & DIJKHUIZEN, L. 2002. *J. Biotechnol.* **94**: 137–155.
- VIGAL, T., GIL, J.A., DAZA, A., GARCIA-GONZALEZ, M.D. & MARTIN, J.F. 1991. *Mol. Gen. Genet.* **225**: 278–288.
- WILSON, J.J. & INGLEDEW, W.M. 1982. *Appl. Environ. Microbiol.* **44**: 301–307.
- WOOD, V., GWILLIAM, R., RAJANDREAM, M.A., LYNE, M. ... NURSE, P. 2002. *Nature* **415**: 871–880.
- YAMAMOTO, T., SHIRAKI, K., FUJIWARA, S., TAKAGI, M., FUKUI, K. & IMANAKA, T. 1999. *Biochem. Biophys. Res. Commun.* **265**: 57–61.
- ZHOU, J., BABA, T., TAKANO, T., KOBAYASHI, S. & ARAI, Y. 1989. *FEBS Lett.* **255**: 37–41.

Received November 22, 2004

Accepted March 16, 2005